The AGAINER Ski **Exoskeleton reduces**

quadriceps muscle

activity by 40%.

Passive Knee Exoskeleton Reduces Quadriceps Muscle Activation During Downhill Skiing: A Pilot Study

Alejandro F. Azocar, Jennifer K. Leestma, Pawel Kudzia, Maria Lazzaroni, Yixing



Liu, Cristina Bayon, Wolfgang F. Rampeltshammer, and Edwin van Asseldonk

INTRODUCTION

- Passive knee exoskeleton with adjustable gas springs
- Assists in maintaining squat-like position through knee extension moment
- Tested effect of exoskeleton assistance on thigh muscle activity during squats and skiing

METHODS

Squat Experiment

- Subject performed squats under the following conditions:
 - \circ No Exo
 - Exo OFF (no assistance)
 - Exo 10 (pressure: 10 bar)

RESULTS



Muscles Tested

- Rectus femoris (RF)
- Vastus lateralis (VL)
- Biceps femoris (BF)
- Semitendinosus (ST)

Electromyography (EMG) Data Processing

- 1. High-pass filtered (20 Hz, 4th order)
- 2. Full-wave rectified
- 3. Low-pass filtered (6 Hz, 2nd order)

- Exo 15 (pressure: 15 bar)
- Exo 20 (pressure: 20 bar)

Ski Experiment

- Subject skied through slalom-style course (requiring 3 turns in each direction):
 - \circ No Exo
 - \circ Exo OFF
 - Exo 15

DISCUSSION

- Quadriceps activity reduced by up to 40%
- Quadriceps activity decreased with increasing exoskeleton pressure
- No effect on hamstrings activity
- Limitations- limited sample size, trials per subject, and testing conditions

DI TECNOLOGIA

4. EMG averaged across trials and normalized by the average EMG from No Exo









This work was performed at the 2019 Winter School on Wearable Robotics, and supported by the European Cooperation in Science and Technology (EU COST) Action Network CA16116.